

11/27/00



JC945 U.S. PTO

11-28-00

Box 898 A

# UTILITY PATENT APPLICATION TRANSMITTAL

Submit an original and a duplicate for fee processing  
(Only for new nonprovisional applications under 37 CFR 1.53(b))

## ADDRESS TO:

Assistant Commissioner for Patents  
Box Patent Application  
Washington, D.C. 20231

Attorney Docket No. 97,017-P7  
First Named Inventor Rodgers  
Express Mail No. EL118813250US  
Total Pages 96



09/22/00

## APPLICATION ELEMENTS

1. ☒ Transmittal Form with Fee
2. ☒ Specification (including claims and abstract)  
[Total Pages 47]
3. ☒ Drawings [Total Sheets 37]
4. ☐ Oath or Declaration [Total Pages ]
  - a. ☐ Newly executed
  - b. ☐ Copy from prior application  
[Note Boxes 5 and 17 below]
    - i. ☐ Deletion of Inventor(s) Signed  
statement attached deleting inventor(s) named  
in the prior application
5. ☐ Incorporation by Reference: The entire disclosure  
of the prior application, from which a copy of the oath or  
declaration is supplied under Box 4b, is considered as  
being part of the disclosure of the accompanying  
application and is hereby incorporated by reference  
therein.
6. ☐ Microfiche Computer Program
7. ☒ Nucleotide and/or Amino Acid Sequence  
Submission
  - a. ☒ Computer Readable Copy
  - b. ☒ Paper Copy
  - c. ☒ Statement verifying above copies

## ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers
9. ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)
  - ☐ PTO-1449 Form
  - ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard  
(Should be specifically itemized)
14. ☒ Small Entity Status Claimed
15. ☐ Certified Copy of Priority Document(s)
16. ☐ Other:

17. ☒ This is a CONTINUING APPLICATION. Please note the following:
- a. ☒ This is a ☐ Continuation ☐ Divisional ☒ Continuation-in-part  
of prior application 09/307,940 filed May 10, 1999
  - b. ☐ Cancel in this application original claims \_\_\_\_\_ of the prior application before calculating the  
filing fee.
  - c. ☒ Amend the specification by inserting before the first line the sentence:  
This is a ☐ continuation ☐ divisional ☒ continuation-in-part  
of application Serial No. 09/307,940 filed May 10, 1999
  - d. ☐ The prior application is assigned of record to

## UTILITY PATENT APPLICATION TRANSMITTAL

Attorney Docket No. 97,017-P7

## APPLICATION FEES

BASIC FEE				\$ 710.00
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	34 -20=	14	x \$18.00	\$ 252.00
Independent Claims	1 - 3=		x \$78.00	\$
<input type="checkbox"/> Multiple Dependent Claims(s) if applicable			+ \$270.00	\$
Total of above calculations =				\$ 962.00
Reduction by 50% for filing by small entity =				\$(481.00)
<input type="checkbox"/> Assignment fee if applicable			+ \$40.00	\$
TOTAL =				\$481.00

18. ☐ Please charge my Deposit Account No. 13-2490 in the amount of \$19. ☒ A check in the amount of \$481.00 is enclosed.

20. The Commissioner is hereby authorized to credit overpayments or charge any additional fees of the following types to Deposit Account No. 13-2490:

- a. ☐ Fees required under 37 CFR 1.16.  
 b. ☐ Fees required under 37 CFR 1.17.  
 c. ☐ Fees required under 37 CFR 1.18.


21. ☐ The Commissioner is hereby generally authorized under 37 CFR 1.136(a)(3) to treat any future reply in this or any related application filed pursuant to 37 CFR 1.53 requiring an extension of time as incorporating a request therefor, and the Commissioner is hereby specifically authorized to charge Deposit Account No. 13-2490 for any fee that may be due in connection with such a request for an extension of time.

## 22. CERTIFICATE OF MAILING

I hereby certify that I directed that the correspondence identified above be deposited with the United States Postal Service as "Express Mail Post Office to Addressee" under 37 CFR § 1.10 on the date indicated below and is addressed to the Asst. Commissioner for Patents, Box Patent Application, Washington, DC 20231.

## 23. USPTO CUSTOMER NUMBER

PATENT &amp; TRADEMARK OFFICE



020306

## 24. CORRESPONDENCE ADDRESS

Name	McDonnell Boehnen Hulbert & Berghoff
Address	300 S. Wacker Drive Suite 3200
City, State, Zip	Chicago, IL 60606

## 25. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

Name Reg. No.	David Harper 42,636
Signature	
Date	11/27/2000

**PATENT**

**APPLICATION FOR PATENT IN THE UNITED STATES PATENT &  
TRADEMARK OFFICE**

**(Case No. 97,017-P7)**

**Title: METHODS FOR TREATING A PATIENT UNDERGOING  
CHEMOTHERAPY**

**Inventors:** Kathleen E. Rodgers  
4403 Galeano Street  
Long Beach, CA 90815

**Who is a citizen of the United States**

Gere diZerega  
1270 Hillcrest Avenue  
Pasadena, CA 91106

**Who is a citizen of the United States**

**Assignee: University of Southern California  
3716 South Hope Street  
Suite 313  
Los Angeles, CA 90007-4344**

**METHODS FOR TREATING A PATIENT UNDERGOING CHEMOTHERAPY**

**Cross Reference**

5           This application claims priority to U.S. Provisional Patent Application Nos. 60/201,470  
filed May 3, 2000; 60/220,804 filed July 25, 2000; 60/233,375 filed September 18, 2000;  
60/235,040 filed September 25, 2000; and 60/243,955 filed October 27, 2000; and is a  
Continuation-In-Part of U.S. Patent Application Serial No. 09/307,940, filed May 10, 1999. This  
Application is related to co-pending and commonly owned patent applications 09/012,400 filed  
10   January 23, 1998; 09/564,051 filed May 3, 2000; 09/564,045 filed May 3, 2000; 09/264,563 filed  
March 8, 2000; 09/245,680 filed February 8, 1999; and 09/658,315 filed September 8, 2000.

**Field of the Invention**

15           The present invention relates to methods, pharmaceutical compositions, and articles of  
manufacture for treating a patient undergoing chemotherapy, particularly for increasing  
hematopoietic cell survival and stem cell mobilization, and reducing the incidence and/or  
severity of chemotherapy-related side effects.

**Background of the Invention**

20           People diagnosed as having cancer are frequently treated with single or multiple  
cytotoxic chemotherapeutic agents (cytotoxic agents) to kill cancer cells at the primary tumor  
site or at distant sites to where cancer has metastasized. (U.S. Patent No. 5,605,931 incorporated  
by reference herein in its entirety.) Chemotherapy treatment is given either in a single or in  
several large doses or, more commonly, it is given in small doses 1 to 4 times a day over variable

times of weeks to months. There are many cytotoxic agents used to treat cancer, and their mechanisms of action are generally poorly understood.

Irrespective of the mechanism, useful chemotherapeutic agents are known to injure and kill cells of both tumors and normal tissues. The successful use of chemotherapeutic agents to treat cancer depends upon the differential killing effect of the agent on cancer cells compared to its side effects on critical normal tissues. Among these effects are the killing of hematopoietic blood forming cells, and the killing and suppression of the white blood cells, which can lead to infection. Acute and chronic bone marrow toxicities are also major limiting factors in the treatment of cancer. They are both related to a decrease in the number of hemopoietic cells (e.g., pluripotent stem cells and other progenitor cells) caused by both a lethal effect of cytotoxic agents or radiation on these cells, and via differentiation of stem cells provoked by a feed-back mechanism induced by the depletion of more mature marrow compartments. (U.S. Patent No. 5,595,973 incorporated by reference herein in its entirety.) Stimulators and inhibitors of bone marrow kinetics play a prominent role in the induction of damage and recovery patterns (Tubiana, M., et al., Radiotherapy and Oncology 29:1, 1993).

Prevention of, or protection from, the side effects of chemotherapy would be a great benefit to cancer patients. The many previous efforts to reduce these side effects have been largely unsuccessful. For life-threatening side effects, efforts have concentrated on altering the dose and schedules of the chemotherapeutic agent to reduce the side effects. Other options are becoming available, such as the use of granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage-CSF (GM-CSF), epidermal growth factor (EGF), interleukin 11, erythropoietin, thrombopoietin, megakaryocyte development and growth factor, pixykinases, stem cell factor, FLT-ligand, as well as interleukins 1, 3, 6, and 7, to increase the number of normal

cells in various tissues before the start of chemotherapy (See Jimenez and Yunis, Cancer Research 52:413-415; 1992). The mechanisms of protection by these factors, while not fully understood, are most likely associated with an increase in the number of normal critical target cells before treatment with cytotoxic agents, and not with increased survival of cells following chemotherapy.

Acute myelosuppression as a consequence of cytotoxic chemotherapy is well recognized as a dose-limiting factor in cancer treatment. (U.S. Patent No. 5,595,973) Although other normal tissues may be adversely affected, bone marrow is particularly sensitive to the proliferation-specific treatment such as chemotherapy or radiotherapy. For some cancer patients, hematopoietic toxicity frequently limits the opportunity for chemotherapy dose escalation. Repeated or high dose cycles of chemotherapy may be responsible for severe stem cell depletion leading to serious long-term hematopoietic sequelae and marrow exhaustion.

Despite advances in the field of chemotherapy, prior art methods have proven to be of limited utility in minimizing chemotherapy-induced depletion of hematopoietic stem cells and their progeny. Thus, there is a need for improved therapeutic methods and pharmaceutical compositions for increasing hematopoietic cell survival following chemotherapeutic treatments, as well as for decreasing the adverse effects of chemotherapy on the bone marrow.

**Summary of the Invention**

In one aspect, the present invention provides methods, pharmaceutical compositions, and articles of manufacture for treating a patient undergoing chemotherapy, for increasing hematopoietic cell survival following chemotherapy, for reducing or preventing other side effects of chemotherapy, such as anemia, and for mobilizing hematopoietic progenitor cells from bone marrow into peripheral blood, comprising administering an amount effective for such purposes

of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof, AII AT<sub>2</sub> type 2 receptor agonists, or ACE inhibitors.

These aspects and other aspects of the invention become apparent in light of the following detailed description.

### **Brief description of the drawings**

**Figure 1** is a graph showing the effect of AII treatment on white blood cell number in the blood 7 days after 5FU treatment.

**Figure 2** is a graph showing the effect of AII treatment on white blood cell number in the spleen 7 days after 5FU treatment.

**Figure 3** is a graph showing the effect of AII treatment on white blood cell number in the thymus 7 days after 5FU treatment.

**Figure 4** is a graph showing the effect of AII treatment on white blood cell number in the bone marrow 7 days after 5FU treatment.

**Figure 5** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following blood harvest 7 days after 5FU treatment.

**Figure 6** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following spleen harvest 7 days after 5FU treatment.

**Figure 7** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following bone marrow harvest 7 days after 5FU treatment.

**Figure 8** is a graph showing the effect of AII treatment on CFU-GM cell number in the blood on day 7 after 5FU treatment.

**Figure 9** is a graph showing the effect of AII treatment on white blood cell number in the spleen on day 14 after 5FU treatment.

**Figure 10** is a graph showing the effect of AII treatment on white blood cell number in the thymus on day 14 after 5FU treatment.

5 **Figure 11** is a graph showing the effect of AII treatment on white blood cell number in the bone marrow on day 14 after 5FU treatment.

**Figure 12** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following spleen harvest 14 days after 5FU treatment.

10 **Figure 13** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following blood harvest 14 days after 5FU treatment.

**Figure 14** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following blood harvest 14 days after 5FU treatment.

**Figure 15** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following spleen harvest 14 days after 5FU treatment.

15 **Figure 16** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following bone marrow harvest 7 days after 5FU treatment.

**Figure 17** is a graph showing the effect of AII treatment on CFU-GM cell number on day 7 after culture initiation following femur harvest 7 days after 5FU treatment.

20 **Figure 18** is a graph showing the effect of AII treatment on CFU-GM cell number in the bone marrow on day 7 after 5FU treatment.

**Figure 19** is a graph showing the effect of AII treatment on CFU-GM cell number in the spleen on day 7 after 5FU treatment.



**Figure 20** is a graph showing the effect of AII treatment on white blood cell number in the blood on day 14 after 5FU treatment.

**Figure 21** is a graph of a different experiment showing the effect of AII treatment on white blood cell number in the blood on days 4, 7, and 10 after 5FU treatment.

5 **Figure 22** is a graph showing the effect of AII(1-7) treatment on white blood cell number in the blood on day 14 after 5FU treatment.

**Figure 23** is a graph showing the effect of 1GD treatment on white blood cell number in the blood on day 14 after 5FU treatment.

10 **Figure 24** is a graph showing the effect of 2GD treatment on white blood cell number in the blood on day 14 after 5FU treatment.

**Figure 25** is a graph showing the effect of 5GD treatment on white blood cell number in the blood on day 14 after 5FU treatment.

**Figure 26** is a graph showing the effect of 9GD treatment on white blood cell number in the blood on day 14 after 5FU treatment.

15 **Figure 27** is a graph showing the effect of 10  $\mu$ g AII and AII analogues and fragments on GM-CFU numbers in the bone marrow on day 10 after 5FU treatment.

**Figure 28** is a graph showing the effect of 100  $\mu$ g AII and AII analogues and fragments on GM-CFU numbers in the bone marrow on day 10 after 5FU treatment.

20 **Figure 29** is a graph showing the effect of 10  $\mu$ g AII and AII analogues and fragments on GM-CFU numbers in the blood on day 10 after 5FU treatment.

**Figure 30** is a graph showing the effect of 100  $\mu$ g AII and AII analogues and fragments on GM-CFU numbers in the blood on day 10 after 5FU treatment.

**Figure 31a and b.** Female C57Bl/6 mice, 6-8 weeks old, were treated with 200 mg/kg cytoxan by intravenous injection. Two days after (panel a) and two days before (panel b) cytoxan injection, subcutaneous administration of AII(1-7) was initiated. Various times after cytoxan injection, the animals were necropsied and peripheral blood harvested. The number of white blood cells was counted by hematocytometer after red blood cell lysis. An asterisk indicates a result significantly different from saline control ( $p \leq 0.05$ ). These data are mean and standard error of 4 animals per group (dose and time point).

**Figure 32a and b.** Female C57Bl/6 mice, 6-8 weeks old, were treated with 200 mg/kg cytoxan by intravenous injection. Two days after and two days before cytoxan injection, subcutaneous administration of AII(1-7) was initiated. Twenty eight after cytoxan injection, the animals were necropsied and bone marrow (panel a) or peripheral blood (panel b) harvested. The number of GM-CFU formed from cells isolated from bone marrow or peripheral blood after red blood cell lysis by culturing in semi solid medium containing recombinant colony stimulating factors was counted. An asterisk indicates a result significantly different from saline control ( $p \leq 0.05$ ). These data are mean and standard error of 3 animals per group (dose and time point).

**Figure 33.** Effect of AII analogues and enalapril on the increase in white blood cells in the peripheral blood after intravenous administration of 5FU.

**Figure 34.** Effect of AII analogues and enalapril on the increase in white blood cells in the peripheral blood after intravenous administration of 5FU.

**Figure 35.** Effect of AII analogues and enalapril on the increase in white blood cells in the peripheral blood after intravenous administration of 5FU.

**Figure 36.** Effect of AII analogues and enalapril on platelet increase after intravenous administration of 5FU.

**Figure 37.** Effect of AII analogues and enalapril on platelet increase after intravenous administration of 5FU.

**Figure 38.** Effect of AII analogues and enalapril on platelet increase after intravenous administration of 5FU.

5 **Figure 39.** Effect of AII analogues and enalapril on hemoglobin increase after intravenous administration of 5FU.

**Figure 40.** Effect of AII analogues and enalapril on hemoglobin increase after intravenous administration of 5FU.

**Figure 41.** Effect of AII analogues and enalapril on hemoglobin increase after intravenous administration of 5FU.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Within this application, unless otherwise stated, the techniques utilized may be found in any of several well-known references such as: *Molecular Cloning: A Laboratory Manual* (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press), *Gene Expression Technology* (Methods in Enzymology, Vol. 185, edited by D. Goeddel, 1991. Academic Press, San Diego, CA), "Guide to Protein Purification" in *Methods in Enzymology* (M.P. Deutscher, ed., (1990) Academic Press, Inc.); *PCR Protocols: A Guide to Methods and Applications* (Innis, et al. 1990. Academic Press, San Diego, CA), *Culture of Animal Cells: A Manual of Basic Technique*, 2<sup>nd</sup> Ed. (R.I. Freshney. 1987. Liss, Inc. New York, NY), *Gene Transfer and Expression Protocols*, pp. 109-128, ed. E.J. Murray, The Humana Press Inc., Clifton, N.J.), and the Ambion 1998 Catalog (Ambion, Austin, TX).

As defined herein the phrase "hematopoietic cells" refers to undifferentiated hematopoietic stem cells, to committed hematopoietic progenitor cells, and to differentiated hematopoietic cells including, but not limited to megakaryocytes, platelets, red blood cells, monocytes, neutrophils, macrophages, and lymphocytes.

As defined herein, "chemotherapy side effects" encompass any deleterious effects suffered as a result of chemotherapy, including but not limited to hematopoietic toxicity, decreased mobilization of hematopoietic progenitor cells from bone marrow into the peripheral blood; anemia, myelosuppression, pancytopenia, thrombocytopenia, neutropenia, lymphopenia, leukopenia, stomatitis, alopecia, headache, and muscle pain.

Unless otherwise indicated, the term "angiotensin converting enzyme inhibitors" or "ACE inhibitors" includes any compound that inhibits the conversion of the decapeptide angiotensin I to angiotensin II, and include but are not limited to alacepril, alatriopril, altiopril calcium, ancovenin, benazepril, benazepril hydrochloride, benazeprilat, benzazepril, benzoylcaptopril, captopril, captopril-cysteine, captopril-glutathione, ceranapril, ceranopril, ceronapril, cilazapril, cilazaprilat, converstatin, delapril, delapril-diacid, enalapril, enalaprilat, enalkiren, enapril, epicaptopril, foroxymithine, fosfenopril, fosenopril, fosenopril sodium, fosinopril, fosinopril sodium, fosinoprilat, fosinoprilic acid, glycopril, hemorphin-4, idapril, imidapril, indolapril, indolaprilat, libenzapril, lisinopril, lyciumin A, lyciumin B, mixanpril, moexipril, moexiprilat, moveltipril, muracein A, muracein B, muracein C, pentopril, perindopril, perindoprilat, pivalopril, pivopril, quinapril, quinapril hydrochloride, quinaprilat, ramipril, ramiprilat, spirapril, spirapril hydrochloride, spiraprilat, spiropril, spiropril hydrochloride, temocapril, temocapril hydrochloride, teprotide, trandolapril, trandolaprilat, utibapril, zabicipril, zabiciprilat, zofenopril and zofenoprilat. (See for example Jackson, et al., Renin and

Angiotensin in Goodman & Gilman's The Pharmacological Basis of Therapeutics, 9th ed., eds. Hardman, et al. (McGraw Hill, 1996); and U.S. Patent No. 5,977,159.)

U.S. Patent No. 5,015,629 to DiZerega (the entire disclosure of which is hereby incorporated by reference) describes a method for increasing the rate of healing of wound tissue, comprising the application to such tissue of angiotensin II (AII) in an amount that is sufficient for said increase. The application of AII to wound tissue significantly increases the rate of wound healing, leading to a more rapid re-epithelialization and tissue repair. The term AII refers to an octapeptide present in humans and other species having the sequence Asp-Arg-Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:1]. The biological formation of angiotensin is initiated by the action of renin on the plasma substrate angiotensinogen (Clouston et al., *Genomics* 2:240-248 (1988); Kageyama et al, *Biochemistry* 23:3603-3609; Ohkubo et al., *Proc. Natl. Acad. Sci.* 80:2196-2200 (1983); each reference hereby incorporated in its entirety). The substance so formed is a decapeptide called angiotensin I (AI) which is converted to AII by the angiotensin converting enzyme (ACE) which removes the C-terminal His-Leu residues from AI [SEQ ID NO: 37]. AII is a known pressor agent and is commercially available.

Studies have shown that AII increases mitogenesis and chemotaxis in cultured cells that are involved in wound repair, and also increases their release of growth factors and extracellular matrices (diZerega, U.S. Patent No. 5,015,629; Dzau et. al., *J. Mol. Cell. Cardiol.* 21:S7 (Supp III) 1989; Berk et. al., *Hypertension* 13:305-14 (1989); Kawahara, et al., *BBRC* 150:52-9 (1988); Naftilan, et al., *J. Clin. Invest.* 83:1419-23 (1989); Taubman et al., *J. Biol. Chem.* 264:526-530 (1989); Nakahara, et al., *BBRC* 184:811-8 (1992); Stouffer and Owens, *Circ. Res.* 70:820 (1992); Wolf, et al., *Am. J. Pathol.* 140:95-107 (1992); Bell and Madri, *Am. J. Pathol.* 137:7-12 (1990). In addition, AII was shown to be angiogenic in rabbit corneal eye and chick chorioallantoic

membrane models (Fernandez, et al., *J. Lab. Clin. Med.* 105:141 (1985); LeNoble, et al., *Eur. J. Pharmacol.* 195:305-6 (1991).

We have previously demonstrated that angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof, and AII AT<sub>2</sub> type 2 receptor agonists are effective in accelerating wound healing and the proliferation of certain cell types, such as hematopoietic stem and lineage specific cells. See, for example, co-pending U.S. Patent Application Serial Nos. 09/012,400, filed January 23, 1998; 09/198,806 filed November 24, 1998; 09/264,563, filed March 8, 2000; 09/287,674, filed April 7, 1999; 09/255,136 filed February 19, 1999; 09/245,680, filed February 8, 1999; 09/250,703 filed February 15, 1999; 09/246,525 filed February 8, 1999; 09/266,293 March 11, 1999; 09/332,582 filed June 14, 1999; 09/373,962 filed August 13, 1999; and 09/352,191 filed July 12, 1999; as well as U.S. Patent Serial Nos. 5,015,629; 5,629,292; 5,716,935; 5,834,432; and 5,955,430; 6,096,709; 6,110,895.

Angiotensin II and its sarcosine analogue have also been used in combination with cytotoxic drugs to induce hypertension in humans and experimental animals undergoing intra-arterial and intraperitoneal chemotherapy. (Taniguchi et al., *J. Nuclear Medicine* 37:1522-1523 (1996); Morita et al., *Am. J. Clin. Oncol.* 15:188-193 (1992); Ohigashi et al., *Hepato-Gastroenterology* 43:338-345 (1996); *Cancer Chemother. Pharmacol.* 39:113-121 (1996); Kuroiwa et al., *Cancer Chemother. Pharmacol.* 35:357-363 (1995); Li et al., *Br. J. Cancer* 67:975-980 (1993); Dworkin et al., *Br. J. Cancer* 76:1205-1210 (1997); Sato et al., *World J. Surg.* 19:836-842 (1995); Mutoh et al., *Urol. Int.* 48:175-180 (1992). In each of these cases, the use of angiotensin II was intended to selectively increase blood flow to the tumor vasculature relative to normal vasculature, thereby increasing the delivery of cytotoxic agent to the tumor.

None of these studies demonstrated or suggested that the use of angiotensin II or its sarcosine analogue would be effective in increasing hematopoietic cell survival, hematopoietic stem cell mobilization into peripheral blood following chemotherapy, or the reduction in chemotherapy side effects.

5       Based on all of the above, it would be unexpected that the use of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, AII, AII analogues, AII fragments or analogues thereof, AII AT<sub>2</sub> type 2 receptor agonists, or ACE inhibitors would be effective in increasing hematopoietic cell survival following chemotherapy, for reducing or preventing other side effects of chemotherapy, such as anemia, and for mobilizing hematopoietic progenitor cells from bone marrow into peripheral blood.

10       A peptide agonist selective for the AT<sub>2</sub> receptor (AII has 100 times higher affinity for AT<sub>2</sub> than AT<sub>1</sub>) has been identified. This peptide is p-aminophenylalanine 6-AII [(p-NH<sub>2</sub>-Phe)6-AII)], Asp-Arg-Val-Tyr-Ile-Xaa-Pro-Phe [SEQ ID NO.36] wherein Xaa is p-NH<sub>2</sub>-Phe (Speth and Kim, BBRC 169:997-1006 (1990). This peptide gave binding characteristics  
15       comparable to AT<sub>2</sub> antagonists in the experimental models tested (Catalioto, et al., *Eur. J. Pharmacol.* 256:93-97 (1994); Bryson, et al., *Eur. J. Pharmacol.* 225:119-127 (1992).

20       The effects of AII receptor and AII receptor antagonists have been examined in two experimental models of vascular injury and repair which suggest that both AII receptor subtypes (AT<sub>1</sub> and AT<sub>2</sub>) play a role in wound healing (Janiak et al., *Hypertension* 20:737-45 (1992); Prescott, et al., *Am. J. Pathol.* 139:1291-1296 (1991); Kauffman, et al., *Life Sci.* 49:223-228 (1991); Viswanathan, et al., *Peptides* 13:783-786 (1992); Kimura, et al., *BBRC* 187:1083-1090 (1992).


Many studies have focused upon AII(1-7) (AII residues 1-7) or other fragments of AII to evaluate their activity. AII(1-7) elicits some, but not the full range of effects elicited by AII. Pfeilschifter, et al., *Eur. J. Pharmacol.* 225:57-62 (1992); Jaiswal, et al., *Hypertension* 19(Supp. II):II-49-II-55 (1992); Edwards and Stack, *J. Pharmacol. Exper. Ther.* 266:506-510 (1993); Jaiswal, et al., *J. Pharmacol. Exper. Ther.* 265:664-673 (1991); Jaiswal, et al., *Hypertension* 17:1115-1120 (1991); Portsi, et a., *Br. J. Pharmacol.* 111:652-654 (1994).

Other data suggests that the AII fragment AII(1-7) acts through a receptor(s) that is distinct from the AT1 and AT2 receptors which modulate AII activity. (Ferrario et al., *J. Am. Soc. Nephrol.* 9:1716-1722 (1998); Iyer et al., *Hypertension* 31:699-705 (1998); Freeman et al., *Hypertension* 28:104 (1996); Ambuhl et al., *Brain Res. Bull.* 35:289 (1994)). Thus, AII(1-7) activity on a particular cell type cannot be predicted based solely on the effect of AII on the same cell type. In fact, there is some evidence that AII(1-7) often opposes the actions of AII. (See, for example, Ferrario et al., *Hypertension* 30:535-541 (1997))

As hereinafter defined, a preferred class of AT2 agonists for use in accordance with the present invention comprises angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, AII, AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists having p-NH-Phe in a position corresponding to a position 6 of AII. In addition to peptide agents, various nonpeptidic agents (e.g., peptidomimetics) having the requisite AT2 agonist activity are further contemplated for use in accordance with the present invention, as well as compounds fused to the active agent to provide some further desired property.



The active AII analogues, fragments of AII and analogues thereof of particular interest in accordance with the present invention comprise a sequence of at least three contiguous amino acids of groups R<sup>1</sup>-R<sup>8</sup> in the sequence of general formula I



5 wherein R<sup>1</sup> is selected from Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

R<sup>2</sup> is selected from Arg, Lys, Ala, Orn, Ser(Ac), Sar, D-Arg and D-Lys;

R<sup>3</sup> is selected from the group consisting of Val, Ala, Leu, Lys, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

10 R<sup>4</sup> is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ser, Ala, homoSer and azaTyr;

R<sup>5</sup> is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

R<sup>6</sup> is selected from the group consisting of His, Arg or 6-NH<sub>2</sub>-Phe;

R<sup>7</sup> is selected from the group consisting of Pro or Ala; and

15 R<sup>8</sup> is selected from the group consisting of Phe, Phe(Br), Ile and Tyr,

excluding sequences including R<sup>4</sup> as a terminal Tyr group.

Particularly preferred combinations for R<sup>1</sup> and R<sup>2</sup> are Asp-Arg, Asp-Lys, Glu-Arg and Glu-Lys.

In alternate embodiments, the active agents comprise a sequence of at least four, five, six, 20 seven, or eight contiguous amino acids of groups R<sup>1</sup>-R<sup>8</sup> in the sequence of general formula I. In a further alternative, the active agents consist essentially of a sequence of at least four, five, six, seven, or eight contiguous amino acids of groups R<sup>1</sup>-R<sup>8</sup> in the sequence of general formula I.

Compounds falling within the category of AT2 agonists useful in the practice of the invention include the AII analogues set forth above subject to the restriction that R<sup>6</sup> is p-NH<sub>2</sub>-Phe. In a further preferred embodiment of all of the aspects of the invention, the active agent comprises a sequence selected from the group consisting of angiotensinogen, SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO: 34; SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, and SEQ ID NO:42.

Particularly preferred embodiments of this class comprise the following sequences: AII (SEQ ID NO:1), AIII or AII(2-8), Arg-Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:2]; AII(3-8), also known as des1-AIII or AIV, Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:3]; AII(1-7), Asp-Arg-Val-Tyr-Ile-His-Pro [SEQ ID NO:4]; AII(2-7). Arg-Val-Tyr-Ile-His-Pro [SEQ ID NO:5]; AII(3-7), Val-Tyr-Ile-His-Pro [SEQ ID NO:6]; AII(5-8), Ile-His-Pro-Phe [SEQ ID NO:7]; AII(1-6), Asp-Arg-Val-Tyr-Ile-His [SEQ ID NO:8]; AII(1-5), Asp-Arg-Val-Tyr-Ile [SEQ ID NO:9]; AII(1-4), Asp-Arg-Val-Tyr [SEQ ID NO:10]; and AII(1-3), Asp-Arg-Val [SEQ ID NO:11]. Other preferred embodiments include: Arg-norLeu-Tyr-Ile-His-Pro-Phe [SEQ ID NO:12] and Arg-Val-Tyr-norLeu-His-Pro-Phe [SEQ ID NO:13]. Still another preferred embodiment encompassed within the scope of the invention is a peptide having the sequence Asp-Arg-Pro-Tyr-Ile-His-Pro-Phe [SEQ ID NO:31]. AII(6-8), His-Pro-Phe [SEQ ID NO:14] and AII(4-8), Tyr-Ile-His-Pro-Phe [SEQ ID NO:15] were also tested and found not to be effective.

In a particularly preferred embodiment, the active agents of the present invention comprise an amino acid sequence of the following general formula:

Asp-Arg-R1-R2-Ile-His-Pro-R3, wherein


R1 is selected from the group consisting of Val, Pro, Lys, Norleu, and Leu;

R2 is selected from the group consisting of Ala, Tyr, and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe or is absent.

In a most particularly preferred embodiment, the active agent comprises a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, and SEQ ID NO:41.

Another class of compounds of particular interest in accordance with the present invention are those comprising a sequence of the general formula II



in which R<sup>2</sup> is selected from the group consisting of H, Arg, Lys, Ala, Orn, Ser(Ac), Sar, D-Arg and D-Lys;

R<sup>3</sup> is selected from the group consisting of Val, Ala, Leu, norLeu, Lys, Ile, Gly, Pro, Aib, Acpc and Tyr;

R<sup>4</sup> is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ser, Ala homoSer and azaTyr;


R<sup>5</sup> is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

R<sup>6</sup> is selected from the group consisting of His, Arg or 6-NH<sub>2</sub>-Phe;

R<sup>7</sup> is selected from the group consisting of Pro or Ala; and

R<sup>8</sup> is selected from the group consisting of Phe, Phe(Br), Ile and Tyr.

A particularly preferred subclass of the compounds of general formula II has the formula



wherein  $R^2$ ,  $R^3$  and  $R^5$  are as previously defined. Particularly preferred is angiotensin III of the formula Arg-Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:2]. Other preferred compounds include peptides having the structures Arg-Val-Tyr-Gly-His-Pro-Phe [SEQ ID NO:17] and Arg-Val-Tyr-Ala-His-Pro-Phe [SEQ ID NO:18]. The fragment AII(4-8) was ineffective in repeated tests; this is believed to be due to the exposed tyrosine on the N-terminus.

In the above formulas, the standard three-letter abbreviations for amino acid residues are employed. In the absence of an indication to the contrary, the L-form of the amino acid is intended. Other residues are abbreviated as follows:

TABLE 1

Abbreviation for Amino Acids

Me <sup>2</sup> Gly	N,N-dimethylglycyl
Bet	1-carboxy-N,N,N-trimethylmethanaminium hydroxide inner salt (betaine)
Suc	Succinyl
Phe(Br)	p-bromo-L-phenylalanyl
azaTyr	aza- $\alpha'$ -homo-L-tyrosyl
Acpc	1-aminocyclopentane carboxylic acid
Aib	2-aminoisobutyric acid
Sar	N-methylglycyl (sarcosine)

It has been suggested that AII and its analogues adopt either a *gamma* or a *beta* turn (Regoli, et al., *Pharmacological Reviews* 26:69 (1974)). In general, it is believed that neutral

side chains in position  $R^3$ ,  $R^5$  and  $R^7$  may be involved in maintaining the appropriate distance between active groups in positions  $R^4$ ,  $R^6$  and  $R^8$  primarily responsible for binding to receptors and/or intrinsic activity. Hydrophobic side chains in positions  $R^3$ ,  $R^5$  and  $R^8$  may also play an important role in the whole conformation of the peptide and/or contribute to the formation of a  
5 hypothetical hydrophobic pocket.

Appropriate side chains on the amino acid in position  $R^2$  may contribute to affinity of the compounds for target receptors and/or play an important role in the conformation of the peptide. For this reason, Arg and Lys are particularly preferred as  $R^2$ .

For purposes of the present invention, it is believed that  $R^3$  may be involved in the formation of linear or nonlinear hydrogen bonds with  $R^5$  (in the gamma turn model) or  $R^6$  (in the beta turn model).  $R^3$  would also participate in the first turn in a beta antiparallel structure (which has also been proposed as a possible structure). In contrast to other positions in general formula I, it appears that beta and gamma branching are equally effective in this position. Moreover, a single hydrogen bond may be sufficient to maintain a relatively stable conformation.  
10 Accordingly,  $R^3$  may suitably be selected from Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr. Lys has also been found to be effective at position  $R^3$ .  
15

With respect to  $R^4$ , conformational analyses have suggested that the side chain in this position (as well as in  $R^3$  and  $R^5$ ) contribute to a hydrophobic cluster believed to be essential for occupation and stimulation of receptors. Thus,  $R^4$  is preferably selected from Tyr, Thr, Tyr  
20  $(PO_3)_2$ , homoSer, Ser and azaTyr. In this position, Tyr is particularly preferred as it may form a hydrogen bond with the receptor site capable of accepting a hydrogen from the phenolic hydroxyl (Regoli, et al. (1974), *supra*). Ala has also been found to be effective at position  $R^4$ .

In position R<sup>5</sup>, an amino acid with a  $\beta$  aliphatic or alicyclic chain is particularly desirable. Therefore, while Gly is suitable in position R<sup>5</sup>, it is preferred that the amino acid in this position be selected from Ile, Ala, Leu, norLeu, Gly and Val.

In the angiotensinogen, AI, AI analogues, AI fragments and analogues thereof, AII analogues, fragments and analogues of fragments of particular interest in accordance with the present invention, R<sup>6</sup> is His, Arg or 6-NH<sub>2</sub>-Phe. The unique properties of the imidazole ring of histidine (e.g., ionization at physiological pH, ability to act as proton donor or acceptor, aromatic character) are believed to contribute to its particular utility as R<sup>6</sup>. For example, conformational models suggest that His may participate in hydrogen bond formation (in the *beta* model) or in the second turn of the antiparallel structure by influencing the orientation of R<sup>7</sup>. Similarly, it is presently considered that R<sup>7</sup> should be Pro in order to provide the most desirable orientation of R<sup>8</sup>. In position R<sup>8</sup>, both a hydrophobic ring and an anionic carboxyl terminal appear to be particularly useful in binding of the analogues of interest to receptors; therefore, Tyr and especially Phe are preferred for purposes of the present invention.

Analogues of particular interest include the following:

TABLE 2

Angiotensin II Analogues

AII Analogue Name	Amino Acid Sequence	Sequence Identifier
Analogue 1	Asp-Arg-Val-Tyr-Val-His-Pro-Phe	SEQ ID NO: 19
Analogue 2	Asn-Arg-Val-Tyr-Val-His-Pro-Phe	SEQ ID NO: 20
Analogue 3	Ala-Pro-Gly-Asp-Arg-Ile-Tyr-Val-His-Pro-Phe	SEQ ID NO: 21
Analogue 4	Glu-Arg-Val-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 22
Analogue 5	Asp-Lys-Val-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 23
Analogue 6	Asp-Arg-Ala-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 24
Analogue 7	Asp-Arg-Val-Thr-Ile-His-Pro-Phe	SEQ ID NO: 25
Analogue 8	Asp-Arg-Val-Tyr-Leu-His-Pro-Phe	SEQ ID NO: 26
Analogue 9	Asp-Arg-Val-Tyr-Ile-Arg-Pro-Phe	SEQ ID NO: 27

Analogue 10	Asp-Arg-Val-Tyr-Ile-His-Ala-Phe	SEQ ID NO: 28
Analogue 11	Asp-Arg-Val-Tyr-Ile-His-Pro-Tyr	SEQ ID NO: 29
Analogue 12	Pro-Arg-Val-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 30
Analogue 13	Asp-Arg-Pro-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 31
Analogue 14	Asp-Arg-Val-Tyr(PO <sub>3</sub> ) <sub>2</sub> -Ile-His-Pro-Phe	SEQ ID NO: 32
Analogue 15	Asp-Arg-norLeu-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 33
Analogue 16	Asp-Arg-Val-Tyr-norLeu-His-Pro-Phe	SEQ ID NO: 34
Analogue 17	Asp-Arg-Val-homoSer-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 35

The polypeptides of the instant invention may be produced by any standard method, including but not limited to recombinant DNA technology and conventional synthetic methods including, but not limited to, those set forth in J. M. Stewart and J. D. Young, *Solid Phase Peptide Synthesis*, 2nd ed., Pierce Chemical Co., Rockford, Ill. (1984) and J. Meienhofer, *Hormonal Proteins and Peptides*, Vol. 2, Academic Press, New York, (1973) for solid phase synthesis and E. Schroder and K. Lubke, *The Peptides*, Vol. 1, Academic Press, New York, (1965) for solution synthesis. The disclosures of the foregoing treatises are incorporated by reference herein.

In general, these methods involve the sequential addition of protected amino acids to a growing peptide chain (U.S. Patent No. 5,693,616, herein incorporated by reference in its entirety). Normally, either the amino or carboxyl group of the first amino acid and any reactive side chain group are protected. This protected amino acid is then either attached to an inert solid support, or utilized in solution, and the next amino acid in the sequence, also suitably protected, is added under conditions amenable to formation of the amide linkage. After all the desired amino acids have been linked in the proper sequence, protecting groups and any solid support are removed to afford the crude polypeptide. The polypeptide is desalted and purified, preferably chromatographically, to yield the final product.

Preferably, peptides are synthesized according to standard solid-phase methodologies, such as may be performed on an Applied Biosystems Model 430A peptide synthesizer (Applied

Biosystems, Foster City, Calif.), according to manufacturer's instructions. Other methods of synthesizing peptides or peptidomimetics, either by solid phase methodologies or in liquid phase, are well known to those skilled in the art. Alternatively, the active agents can be prepared by standard recombinant DNA techniques.

5 In one aspect, the present invention provides methods and kits for increasing hematopoietic cell survival following chemotherapy, and treating and preventing the adverse effects of chemotherapy, comprising the administration of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists (hereinafter referred to as  
10 "active agents") to a patient undergoing chemotherapy.

In another aspect, the present invention provides methods and kits for mobilizing hematopoietic progenitor cells from bone marrow into peripheral blood comprising the administration of the active agents of the invention to a patient in need of such treatment. This aspect of the invention can also be used to treat a patient in need of chemotherapy.

15 The methods of the invention are appropriate for use with chemotherapy using any cytotoxic agent, including, but not limited to, cyclophosphamide, taxol, 5-fluorouracil, adriamycin, cisplatin, methotrexate, cytosine arabinoside, mitomycin C, prednisone, vindesine, carbaplatin, and vincristine. The cytotoxic agent can also be an antiviral compound which is capable of destroying proliferating cells. For a general discussion of cytotoxic agents  
20 used in chemotherapy, see Sathe, M. et al., Cancer Chemotherapeutic Agents: Handbook of Clinical Data (1978), hereby incorporated by reference.

The methods of the invention are also particularly suitable for those patients in need of repeated or high doses of chemotherapy. For some cancer patients, hematopoietic toxicity



frequently limits the opportunity for chemotherapy dose escalation. Repeated or high dose cycles of chemotherapy may be responsible for severe stem cell depletion leading to severe long-term hematopoietic sequelae and marrow exhaustion. The methods of the present invention provide for improved mortality and blood cell count when used in conjunction with  
5 chemotherapy.

The active agents may be administered by any suitable route, including orally, parentally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles. The term parenteral as used herein includes, subcutaneous, intravenous, intraarterial, intramuscular, intrasternal, intratendinous, intraspinal, intracranial, intrathoracic, infusion techniques, or intraperitoneally.  
10

The active agents may be made up in a solid form (including granules, powders or suppositories) or in a liquid form (*e.g.*, solutions, suspensions, or emulsions). The compounds of the invention may be applied in a variety of solutions. Suitable solutions for use in accordance with the invention are sterile, dissolve sufficient amounts of the peptide, and are not harmful for  
15 the proposed application. In this regard, the compounds of the present invention are very stable but are hydrolyzed by strong acids and bases. The compounds of the present invention are soluble in organic solvents and in aqueous solutions at pH 5-8.

The active agents may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional adjuvants, such as preservatives, stabilizers,  
20 wetting agents, emulsifiers, buffers etc.

For administration, the active agents are ordinarily combined with one or more adjuvants appropriate for the indicated route of administration. The compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, stearic acid, talc, magnesium

stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulphuric acids, acacia, gelatin, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and tableted or encapsulated for conventional administration. Alternatively, the compounds of this invention may be dissolved in saline, water, polyethylene glycol, propylene glycol, carboxymethyl cellulose colloidal solutions, ethanol, corn oil, peanut oil, cottonseed oil, sesame oil, tragacanth gum, and/or various buffers. Other adjuvants and modes of administration are well known in the pharmaceutical art. The carrier or diluent may include time delay material, such as glyceryl monostearate or glyceryl distearate alone or with a wax, or other materials well known in the art.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (*e.g.*, liniments, lotions, ointments, creams, or pastes) and drops suitable for administration to the eye, ear, or nose.

The dosage regimen of active agents for the methods of the invention is based on a variety of factors, including the type of injury, the age, weight, sex, medical condition of the individual, the severity of the condition, the route of administration, and the particular compound employed. Dosage levels of the order of between 0.1 ng/kg and 10 mg/kg body weight of the active agents per body weight are useful for all methods of use disclosed herein.

In all of these embodiments, the compounds of the invention can be administered prior to, simultaneously with, or subsequent to chemotherapeutic exposure.

In a preferred embodiment, the active agent is administered subcutaneously. A suitable subcutaneous dose of the active agent is preferably between about 0.1 ng/kg and about 10 mg/kg administered twice daily for a time sufficient to increase white blood cell survival after chemotherapy treatment or to mobilize hematopoietic progenitor cells from bone marrow into peripheral blood. In a more preferred embodiment, the concentration of active agent is between

about 100 ng/kg body weight and about 10.0 mg/kg body weight. In a most preferred embodiment, the concentration of active agent is between about 2.5 µg/kg body weight and about 100 µg/kg body weight. This dosage regimen maximizes the therapeutic benefits of the subject invention while minimizing the amount of agonist needed. Such an application minimizes costs as well as possible deleterious side effects. For example, the active agents are administered to an oncology patient for up to 30 days prior to a course of chemotherapy and for up to 60 days post-chemotherapy. The therapy is administered for 1 to 6 times per day at dosages as described above. In a further preferred embodiment, the active agent is administered once per day.

For subcutaneous administration, the active ingredient may comprise from 0.0001% to 10% w/w, *e.g.*, from 1% to 2% by weight of the formulation, although it may comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation.

In a preferred embodiment, subcutaneous administration of between about 1 to 1000 µg/kg/day of the active agents is initiated at between one week before to one week after administration of a chemotherapeutic agent. In a most preferred embodiment, administration of the active agents begins either at the time chemotherapy is initiated, or 0-10 days after initiation.

In another preferred embodiment of the invention, a subject undergoes repeated cycles of treatment according to the method of this invention. Preferably, a subsequent treatment cycle commences only after administration of the compounds of the invention has terminated, and the subject's blood cell counts (*e.g.*, white blood cell count, as well as platelet and megakaryocyte count) have returned to a therapeutically acceptable level (as determined by the attending veterinarian or physician), permitting the repeated chemotherapy.

002211260  
In a further aspect, the present invention provides kits for increasing hematopoietic cell survival following chemotherapy, reducing the incidence and/or severity of anemia and other side effects of chemotherapy, and/or mobilizing hematopoietic progenitor cells from bone marrow into peripheral blood, wherein the kits comprise an effective amount of the active agents  
5 for increasing hematopoietic cell survival following chemotherapy, reducing chemotherapy-induced side effects, or for mobilizing hematopoietic progenitor cells from bone marrow into peripheral blood, and instructions for using the amount effective of active agent as a therapeutic. In a preferred embodiment, the kit further comprises a pharmaceutically acceptable carrier, such as those adjuvants described above. In another preferred embodiment, the kit further comprises a  
10 means for delivery of the active agent to a patient. Such devices include, but are not limited to syringes, matrical or micellar solutions, bandages, wound dressings, aerosol sprays, lipid foams, transdermal patches, topical administrative agents, polyethylene glycol polymers, carboxymethyl cellulose preparations, crystalloid preparations (e.g., saline, Ringer's lactate solution, phosphate-buffered saline, etc.), viscoelastics, polyethylene glycols, and polypropylene glycols. The means  
15 for delivery may either contain the effective amount of the active agents, or may be separate from the compounds, which are then applied to the means for delivery at the time of use.

In a further embodiment, the present invention provides an article of manufacture, comprising the pharmaceutical composition of the invention preloaded into a syringe or other delivery system, for home use by patients undergoing chemotherapy.

20 In another aspect, the present invention provides a pharmaceutical composition, comprising an amount effective of the active agents to increase hematopoietic cell survival and/or to reduce the side effects of chemotherapy in a chemotherapy patient, and a

pharmaceutically acceptable carrier. In a preferred embodiment, the active agent is AII(1-7), and the effective dosage is between about 2.5 µg/kg/day and 100 µg/kg/day.

In a further embodiment, the pharmaceutical composition further comprises an amount of cytokine effective for increasing the production of hematopoietic cells. According to this aspect of the invention, cytokines appropriate for use include, but are not limited to, granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage-CSF (GM-CSF), epidermal growth factor (EGF), interleukin 11, erythropoietin, thrombopoietin, megakaryocyte development and growth factor, pixykinines, stem cell factor, FLT-ligand, as well as interleukins 1, 3, 6, and 7. In a most preferred embodiment, the cytokine is granulocyte colony stimulating factor.

The methods, kits, and pharmaceutical compositions of the present invention, significantly enhance the utility of presently available treatments for clinical chemotherapeutic treatments, by providing improved methods for increasing hematopoietic blood cell survival following chemotherapy, reducing the side effects of chemotherapy, and mobilizing hematopoietic progenitor cells from bone marrow into peripheral blood, and also by their cytokine sparing effect, in that significantly reduced amounts of cytokines are needed when administered with the active agents to a patient undergoing chemotherapy.

The present invention may be better understood with reference to the accompanying example that is intended for purposes of illustration only and should not be construed to limit the scope of the invention, as defined by the claims appended hereto.

*Example 1. Effect of AII on white blood cell mobilization and recovery after 5 fluorouracil treatment*

This study was designed to test the effect of AII on the recovery of white blood cells in lymphoid organs and on the level of granulocyte macrophage precursors (CFU-GM) in the blood

(ie: mobilization), spleen (mobilization), and bone marrow (recovery) after intravenous administration of 5-fluorouracil (5FU).

Subcutaneous administration of AII (either 10 or 100 µg/kg/day) was initiated either two days before (-d2), the day of (d0), or two days after (d2) intravenous administration of 5FU. On either day 7 or 14 after 5FU administration, the animals were necropsied and the spleen, thymus, peripheral blood, and bone marrow were harvested. The number of white blood cells in each of the lymphoid organs, or the number of CFU-GM present in all the organs except the thymus, were then assessed. The number of white blood cells per lymphoid organ was assessed after (1) dissociation of the tissues into a single cell suspension (thymus and spleen), (2) flushing of bone marrow from the femur, or (3) lysis of red blood cells by a hypotonic ammonium chloride solution (blood). An aliquot of the cell suspension was diluted with 0.04% trypan blue and the number of cells was determined by microscopic analysis using a hemacytometer. After counting, the number of cells were adjusted to allow a 1:10 dilution of cells into semi-solid medium containing fetal bovine serum, bovine serum albumin, methyl cellulose, stem cell factor, interleukin 3, interleukin 6, L-glutamine, 2 mercaptoethanol, human transferrin and bovine insulin. On day 7 after culture initiation, the number of CFU-GM per well (and then per organ) was determined by microscopic analysis (**Figures 1-20**). These data demonstrate that AII treatment after chemotherapy leads to significantly enhanced white blood cell mobilization and/or recovery in all of the tissues tested.

***Example 2. Effect of AII analogues and fragments on white blood cell mobilization and recovery after 5 fluorouracil treatment***

The method was conducted as described above in Example 1, except that mice were injected subcutaneously with 150 mg/kg body weight of 5FU, and AII peptide analogues and

fragments were tested. Administration of the peptides (see Table 3) was begun 2 days after and continued until 10 days after 5FU administration, at which time the mice were euthenized for evaluation of bone marrow and blood GM-CFU progenitors. On days 4 and 7 after 5FU administration, blood was taken under anesthesia from the retro-orbital sinus. On day 10, blood was taken by cardiac puncture.

The data for these experiments is shown in Figures 21-30, and demonstrate that all peptides tested accelerated the recovery of white blood cells after chemotherapy (Figures 21-26), increased the number of GM-CFU progenitors in the bone marrow (Figures 27-28), and increased the mobilization of GM-CFU progenitors from the bone marrow into the peripheral blood (Figures 29-30), relative to controls. The peptides were effective at both concentrations tested (10 µg/kg/day and 100 µg/kg/day), and the efficacy generally increased with increasing length of treatment.

**Table 3:      Designation for Analogues/Fragments**

Name	Abbreviation	Sequence	SEQ ID NO:
1GD	Ala4-AII(1-7)	DRVAIHP	SEQ ID NO:38
2GD	Pro3-AII(1-7)	DRPYIHP	SEQ ID NO:39
5GD	Lys3-AII(1-7)	DRKYIHP	SEQ ID NO:40
9GD	NorLeu-AII(1-7)	DR(nor)YIHP	SEQ ID NO:41
AII(1-7)		DRVYIHP--	SEQ ID NO:4
AII		DRVYIHPF	SEQ ID NO. 1

**Example 3.      Effect of AII(1-7) on Hematopoietic Recovery After Cytosan**

In this study, female C57Bl/6 mice were injected with 200 mg/kg cyclophosphamide (“cytosan”) intravenously. Administration of AII(1-7) by subcutaneous injection was begun 2 days before or 2 days after administration of the antineoplastic and continued daily until necropsy for evaluation of mature formed blood elements in the circulation and the number of GM-CFU in the bone marrow and peripheral blood. On days 5, 9, 14, 21 and 28 after cytosan

administration, blood was taken under metofane anesthesia from the retro-orbital sinus to assess white blood cell (WBC) and platelet number.

The data showed that AII(1-7) accelerated the recovery of WBC after intravenous administration of cyclophosphamide when treatment with peptides was initiated two days after the chemotherapeutic (Figure 31a). The increase in WBC number was observed within 9 days after cyclophosphamide treatment. However, if AII(1-7) treatment was initiated two days prior to the administration of cyclophosphamide, there was a decrease in WBC number compared with control starting on day 14 after exposure to the chemotherapeutic (Figure 31b).

Administration of cyclophosphamide also significantly reduced the number of myeloid progenitors (GM-CFU) in the bone marrow. Treatment with AII(1-7) (100 µg/kg/day) 2 days prior to treatment with cytoxan slightly reduced the number of GM-CFU in the bone marrow of treated animals (Figure 32a). However, initiation of AII(1-7) administration 2 days after chemotherapy increased the number of myeloid progenitor cells in the bone marrow (Figure 32a). Further, administration of AII(1-7), initiated after cyclophosphamide treatment, increased the number of myeloid progenitors in the peripheral blood (Figure 32b).

#### **Example 4. *Effect of Angiotensin Fragments and Ace Inhibitor on Recovery After Chemotherapy***

The next study was designed to test the effect of angiotensin peptides and the angiotensin converting enzyme inhibitor, enalapril on recovery of white blood cells, platelets and hemoglobin after chemotherapy, as well as on granulocyte-macrophage colony forming units (GM-CFU) in the blood and bone marrow.

Female C67Bl/6 mice (4 per group), 6-8 weeks old, were injected with 150 mg/kg 5FU intravenously. Administration of the peptides (AII (100 µg/kg/day), AII(1-7) (10 or 100



µg/kg/day), AII(1-5) (10 or 100 µg/kg/day), and AII(1-6) (100 µg/kg/day)), by subcutaneous injection, or enalapril (30 mg/kg/day) by oral gavage, was begun 2 days after administration of the antineoplastic and continued daily until 28 days after 5FU administration, at which time the mice were euthanized for evaluation of white blood cell numbers, platelets, hemoglobin levels and the number of GM-CFU in the bone marrow and peripheral blood. On days 7, 10, 14 and 21 after 5FU administration, blood was taken under anesthesia from the retro-orbital sinus to assess white blood cell number, platelet number, and hemoglobin levels.

#### *Retro-orbital Bleeding of Mice*

The mice were bled from the retro-orbital sinus at days 7, 10, 14 and 21. The mice were anesthetized with Metofane (an inhaled anesthesia). Approximately 150-200 µl of blood were obtained from the retro-orbital sinus with a heparinized capillary tube. The blood was then placed in a 1.7 ml microfuge tube containing 10 mM EDTA and held on ice until further processing.

#### *Hemoglobin Assay*

One hundred µl of blood was pipetted into a centrifuge tube, to which 900 µl of distilled water was added. The blood and water were mixed by inversion and allowed incubate at 4°C for 20 minutes. The tube was then centrifuged at 12000 rpm to precipitate the cellular debris for 20 to 30 minutes at room temperature. Twenty µl of the supernatant from this centrifugation was then added into triplicate wells of a 96 well plate containing 180 µl of distilled water. The optical density was then read using a microplate reader at 570 nM.

#### *WBC and Platelet Evaluation*

Twenty  $\mu$ l of blood was mixed with 200  $\mu$ l of red blood cell (RBC) lysing solution (0.83%  $\text{NH}_4\text{Cl}$ , 10 mM EDTA, 0.5%  $\text{NaHCO}_3$ ). The mixture was then incubated for 10 minutes at 4°C. After this incubation, the supernatant was removed and the pellet was resuspended in 100  $\mu$ l of PBS. To this, 100  $\mu$ l of 0.04% trypan blue was added. This mixture was vortexed and the number of WBC (baseline was approximately  $10^7$  cells/ml) was evaluated by hematocytometer under light microscopy and the number of platelets (baseline was approximately  $2.5 \times 10^8$  platelets/ml) was evaluated by hematocytometer under phase contrast microscopy.

#### *Evaluation of GM-CFU Progenitors in the Blood and Bone Marrow*

The blood was harvested by cardiac puncture on day 10 to assess mobilization of myeloid progenitors into the peripheral blood. The femurs and tibia were also collected and the bone marrow was harvested by flushing with PBS containing 2% fetal calf serum. After collection of the blood and bone marrow, the red blood cells were lysed with a hypotonic solution (described above), mixed with 0.04% trypan blue, and the number of nucleated cells assessed by hematocytometer under light microscopy. Aliquots of cells were then resuspended at  $1 \times 10^5$  cells/ml (bone marrow) or  $1 \times 10^6$  cells/ml (blood). One hundred  $\mu$ l of each suspension was added to 900  $\mu$ l of semisolid medium containing 0.9% methyl cellulose in Iscove's MDM, 15% fetal calf serum, 1% bovine serum albumin, 10  $\mu$ g/ml bovine pancreatic insulin, 200  $\mu$ g/ml human transferrin,  $10^{-4}$  M 2-mercaptoethanol, 2 mM glutamine, 10 ng/ml recombinant murine interleukin 3, 10 ng/ml recombinant human interleukin 6, 50 ng/ml recombinant murine stem cell factor and 3 units/ml erythropoietin. This mixture was then added to duplicate wells of a 24 well plate. The cultures were then placed at 37°C in a humidified atmosphere of 5%  $\text{CO}_2$  in air. At

day 14, the number of myeloid colonies formed was enumerated under phase contrast microscopy.

### *Results*

These studies were conducted to compare the effect of angiotensin fragments on hematopoietic recovery after chemotherapy. No animals were lost to analysis as a result of these therapies. The animals that died succumbed to anesthesia during the bleeding procedures.

There was no difference in baseline white blood cell (WBC) number between groups (8.6 to  $9.0 \times 10^6$  per ml). Baseline platelet numbers ranged from  $2.4$  to  $2.6 \times 10^8$  platelets per ml. No differences were observed between the groups at baseline.

These studies showed that all of the active agents tested increased the number of white blood cells in the peripheral blood after intravenous administration of 5FU (Figures 33-35). The decrease in WBC number as a result of administration of 5FU reached approximately 84%. The increase in WBC number was observed within 7 days after 5FU treatment (within 5 days after initiation of peptide administration). The increase in WBC continued throughout the experimental period but varied with the active agent. AII led to the most profound effect on this parameter followed by AII(1-7) and AII(1-5). AII(1-6) and enalapril were the least effective.

Further, there was also an increase in platelets after administration of both AII(1-7) and AII(1-5) (Figures 36-38). Intravenous administration 5FU resulted in approximately a 70% decrease in platelet number that was maximal at day 10. Increases in platelet number occurred on days 14 and 21 with all test articles increasing the concentration of platelets in the blood.

An increase in the level of hemoglobin in the blood was observed with all active agents, except AII, on day 10 (the nadir in control animals in this parameter). Thereafter, no further effect was observed (Figures 39-41).

In summary, all active agents tested accelerated the recovery of white blood cells, platelet number, and hemoglobin level after chemotherapy.

**Example 5.** *Phase I/II Dose Escalation Study of AII(1-7) (SEQ ID NO:4) Administered Before and After Chemotherapy in Patients with Newly Diagnosed Breast Cancer*

Delivery of optimal dosing of cytotoxic chemotherapy is often limited by myelosuppression. Erythropoietin, filgrastim (G-CSF), sargramostim (GM-CSF), and oprelvekin (IL-11) have been the first recombinant hematopoietic growth factors to be United States Food and Drug Administration approved to stimulate human blood production and mitigate the toxicities of cytotoxic chemotherapy. There are a number of additional hematopoietic regulatory molecules that have been identified and are being produced in sufficient quantities to permit clinical testing in humans.

Data derived from pre-clinical studies demonstrated the effectiveness of AII(1-7) (SEQ ID NO:4) in accelerating hematopoietic recovery following chemotherapy induced myelosuppression. Additional activity on bone marrow and peripheral blood progenitor mobilization and proliferation was demonstrated with AII(1-7) indicating a potential for clinical utility following myelosuppressive cancer therapies. The pharmacologic effects appear to be multi-lineage and dose dependent.

The hematopoietic properties demonstrated in the pre-clinical studies support the investigation into the usefulness of AII(1-7) to decrease the incidence and severity of complications associated with myelosuppression secondary to cytotoxic therapy.

Based upon safety evaluation studies in animals (where the daily doses tested ranged from 10 to 1,000 µg/kg for 30 days) and the pharmacology profile of AII(1-7), the dose range

used in humans ranged from 2.5 to 100 µg/kg/day over at least 10 days. This provided for an approximate 9-10 fold safety margin over the animal exposures.

### **Study Objectives**

#### **Primary objectives**

- 5 a) Determine the optimal biologic dose (OBD) or maximum tolerated dose (MTD) of AII(1-7) in cancer patients before and after treatment with cytotoxic therapy.

#### **Secondary objectives**

- 10 a) Assess the hematologic profile in time to nadir, nadir, and time to recovery (absolute neutrophil count (ANC) > 500 cells/µL and platelets > 25,000/µL) in patients treated with AII(1-7) after treatment with chemotherapy.
- b) Assess the mobilization of CD34+ progenitor cells and colony forming units (CFU-GM and CFU-GEMM) in peripheral blood after AII(1-7) treatment given before and after treatment with cytotoxic chemotherapy.
- 15 c) Assess the pharmacokinetic profile of AII(1-7) treatment given before treatment with cytotoxic therapy.
- d) Assess the incidence and days of hospitalization, febrile neutropenia ( $\geq 38.2^{\circ}$  C and ANC < 1000/µL), and days of antibiotic use compared to filgrastim.
- e) Assess the influence of AII(1-7) on the chemotherapy regimen (disease free survival (DFS) and overall survival (OS)).
- 20 f) Assess any synergy in hematologic response with AII(1-7) in combination with filgrastim.

### **Investigational Plan**

25 This study compared the effects of AII(1-7) (SEQ ID NO:4) in patients with newly diagnosed breast cancer receiving doxorubicin 60 mg/m<sup>2</sup> and cyclophosphamide 600 mg/m<sup>2</sup> for at least 3 cycles of adjuvant chemotherapy following surgical tumor reduction. A filgrastim (recombinant G-CSF) (NEUPOGEN®, Amgen, Inc., Thousand Oaks, CA) comparator arm was used to compare safety and response variables and to assess synergy of AII(1-7) (SEQ ID NO:4)

30 with filgrastim.

### *Dose Escalation Scheme*

Patients who satisfied the inclusion/exclusion criteria received a once daily subcutaneous injection of the given AII(1-7) dose level for 7 days followed by a 1 week rest period prior to any chemotherapy (cycle 0), in order to permit evaluation of potential side effects of AII(1-7) in the absence of toxicity due to chemotherapy. Dose escalation within an individual patient was not be permitted. Dose escalation proceeded based on the occurrence of dose limiting toxicity (DLT).

### *Post Chemotherapy Studies*

Following a rest period of 7 days, a chemotherapy regimen containing doxorubicin 60 mg/m<sup>2</sup> and cyclophosphamide 600 mg/m<sup>2</sup> was initiated. AII(1-7) was administered for at least 10 days, or until the ANC > 1500/μL for 2 days, beginning two days after chemotherapy. Up to three chemotherapy cycles followed by AII(1-7) administration were repeated every 21 days or as indicated by patient tolerance. Any patient that failed to achieve an ANC > 1500/μL by day 15 (13 days of AII(1-7)) received filgrastim at 5.0 μg/kg/day until the ANC > 1500/μL for 2 days. Dosing began with the lowest dose of AII(1-7). In combination with each dosing group of AII(1-7), one additional patient received filgrastim 5.0 μg/kg/day beginning two days after chemotherapy and continuing for at least 10 days or until the ANC > 1500/μL for 2 days. These patients were used for comparison with the AII(1-7) treated patients. Any patient in the AII(1-7) dosing arm that experienced an episode of febrile neutropenia following chemotherapy or failed to achieve an ANC > 1500/μL during the first cycle received filgrastim at 5.0 μg/kg/day in combination with AII(1-7) for the remaining chemotherapy cycles. The remaining cycles for that patient were treated with filgrastim off protocol as determined by the Investigator. Patients received supportive treatment consistent with the standard of care as determined by the Investigator.

### *Endpoints*

The primary safety endpoints were the incidence and grade of toxicity experienced by each dose group (DLT), MTD or OBD, changes in biochemistry, hematology, urinalysis, physical findings, and adverse events. Secondary safety endpoints were DFS and OS.

The preliminary efficacy endpoints were time to nadir (i.e.: the low point), nadir, and hematologic recovery (ANC > 500/ $\mu$ L and platelets > 25,000/ $\mu$ L), mobilization of CD34+ progenitor cells, CFU-GM and GEMM after AII(1-7) (SEQ ID NO:4) treatment given before and after cytotoxic chemotherapy, and incidence and number of days of febrile neutropenia ( $\geq 38.2^{\circ}$  C; ANC < 1000/ $\mu$ L), hospitalization, infection, and antibiotic use.

## Treatments

AII(1-7), filgrastim or both were self-administered following adequate patient training. AII(1-7) was administered once daily with a subcutaneous needle into a site located in the abdomen or thigh every morning.

## Efficacy and Safety Variables

### *Hematology*

Hematology assessment included red blood cell count (RBC), hemoglobin (Hgb), hematocrit (Hct), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), white blood cell count (WBC) including differential, reticulocyte count, and platelet count.

### *Coagulation*

A coagulation panel was obtained for prothrombin time (PT) and activated partial thromboplastin time (PTT).

### *Adverse Event*

An adverse event (AE) was considered as any unfavorable or unintended change in structure, function, signs, or symptoms temporally associated with the use of a medicinal product experienced by a person administered a pharmaceutical product, whether or not a causal relationship with the product was established. Clinically significant laboratory abnormalities were considered AEs if deemed appropriate by the Investigator. Worsening of a pre-existing condition was also considered an AE, as was the discovery of an abnormal finding during physical exam that was not included in the medical history.

### *Appropriateness of Measurements*

The occurrence of febrile neutropenia was associated with significant morbidity and mortality in patients following myelosuppressive chemotherapy.

### *Primary Efficacy Variables*

#### *CD34+ assays*

Peripheral blood was collected to conduct CD 34+ cell mobilization assays. Blood samples were obtained with a draw volume of 2 mL blood collection tubes.

#### *Febrile Neutropenia*

Patients were instructed to notify the Investigator of any oral temperature  $\geq 38.2^{\circ}\text{C}$ . Patients had a CBC obtained anytime they present with fever. If the ANC was  $< 1000/\mu\text{L}$  in conjunction with an oral temperature  $\geq 38.2^{\circ}\text{C}$ , two blood cultures for aerobic/anaerobic bacteria were obtained, drawn 15 minutes apart.

#### *Drug Concentration Measurements*

Pharmacokinetics samples for the measurement of AII(1-7) and AII were collected and centrifuged. The specimen was decanted into a plastic transfer vial and stored at  $-20^{\circ}\text{C}$ . Blood samples were obtained with a draw volume of 5 mL.

### **Results:**

Administration of AII(1-7) resulted in an increase in multiple hematopoietic lineages and mobilization of hematopoietic progenitors into the peripheral blood. A dose dependent increase in the level of nadir in the number of white blood cells and absolute neutrophil count and time to these nadirs was observed. This is consistent with an effect on the compound on myeloid recovery after chemotherapy. At the lower doses, administration of filgrastim was required on Day 15 to restore acceptable levels of WBC prior to the next chemotherapy cycle. However, only one to two doses were required to substantially increase WBC and ANC suggesting a



priming of the bone marrow cells for response to differentiating colony stimulating factors and a cytokine sparing effect of AII(1-7).

An additional benefit of administration of AII(1-7) was the ability of all patients to maintain on cycle, full intensity chemotherapy. It is expected that, after the first cycle of chemotherapy, the full dose of chemotherapeutic drug would not be administered to the patient at the time that would be optimal for cancer therapy, due to a dose-limiting toxicity that requires resolution. However, AII(1-7) administration permitted the patients to maintain full intensity chemotherapy on cycle.

Further, administration of AII(1-7) resulted in a dose-dependent increase in platelets. At the lowest dose, the nadir in platelet number occurred on day 12 and platelet number recovered thereafter. At 50 µg/kg/day of AII(1-7), only day 12 showed any change in platelet number. At the next higher dose, no decrease in platelet number occurred. In fact, an early increase relative to baseline was observed. In contrast, patients that received filgrastim had a time dependent decrease in platelet number that was more pronounced in subsequent cycles of chemotherapy.

Administration of AII(1-7) also affected the correction of anemia following administration of chemotherapeutic drugs. It was expected that administration of the chemotherapeutic drug would reduce the blood hemoglobin levels, that this reduction would not be corrected within the chemotherapy cycle and that the anemia would get progressively worse. This was observed after administration of filgrastim in these patients. After administration of the lowest dose of AII(1-7), a reduction in hemoglobin was observed. Contrary to that expected after administration of the chemotherapeutic drug in the absence of an adjuvant, the hemoglobin level returned to baseline levels by the next cycle. At subsequent dose increases of AII(1-7), slight to no anemia was observed, but at all doses, restoration of hemoglobin prior to the next cycle was observed.

To date, 14 patients have been exposed to AII(1-7) at 4 dosages (2.5 µg/kg/d, 10 µg/kg/d, 50 µg/kg/d, 75 µg/kg/d, and 100 µg/kg/d). Cumulative doses of >1500 µg/kg have been reached. No acute affect on blood pressure following administration has been observed in patients with or without a history of hypertension. No dose-limiting toxicities have been observed with AII(1-7) and all study patients have received 100% dose intensive chemotherapy to date. Anemia correction with each cycle has been observed with doses as low at 2.5 µg/kg. No drug related serious adverse events have been observed. No cycle delays due to neutropenia have occurred,

however, patients treated at 2.5 and 10 µg/kg have needed 1-2 doses of filgrastim at day 15 to treat grade 4 neutropenia as directed by the protocol. Of note is that following 10-12 days of AII(1-7), as little as 1 dose of filgrastim normalizes the neutrophil count indicating a possible synergy between filgrastim and AII(1-7).

AII(1-7) reduced the frequency of grade 2-4 thrombocytopenia, grade 2-4 anemia, and grade 3-4 lymphopenia compared to filgrastim. Filgrastim patients experienced a lower frequency of grade 3-4 neutropenia compared to AII(1-7), however, the frequency of grade 3-4 leukopenia was similar in both groups. The most prominent hematologic effect with AII(1-7) was the prevention of thrombocytopenia.

Toxicity	Filgrastim	2.5 µg/kg	10 µg/kg	50 µg/kg	75 µg/kg	100 µg/kg
Hgb <10gm/dl	60%	66%	33%	0%	0%	0%
Platelet <80K/µl	60%	0%	0%	0%	0%	0%
Lymph <500/µl	80%	66%	33%	33%	0%	0%
ANC <1000/µl	40%	100%	100%	100%	100%	100%
WBC <2000/µl	60%	66%	66%	66%	100%	33%

Hgb= hemoglobin

Lymph= lymphocyte

ANC= Absolute neutrophil count

WBC= white blood cells

Additionally, AII(1-7) reduced the frequency of stomatitis by 30% of that observed with filgrastim treatment, as well as decreasing the frequency of a number of other common side effects of chemotherapy, such as headache, muscle pain, and alopecia, relative to both historical numbers and filgrastim controls.

Treatment Group	Historic	AII(1-7) (n=15)	Filgrastim (n=5)
Nausea	90%	93%	100%
Fatigue		93%	80%
Anemia	22%	7%	60%
Vomiting		80%	60%
Headache		26%	60%
Stomatitis	88%	40%	60%
Myalgia/Muscle-skeletal pain		53%	60%
Alopecia	77%	46%	40%

In summary, the data demonstrate the following:

-AII(1-7) is safe without observed dose limiting toxicities.

-Doses of AII(1-7) greater than 10 µg/kg/day appear to be active; the optimal dosage range appears to be 50-75 µg/kg/day.

-No over-production of formed blood elements occurs before or after chemotherapy.

-AII(1-7) reduces the frequency of gradable anemia, lymphopenia, and thrombocytopenia compared to filgrastim treated patients.


-Neutrophil nadirs are not affected by AII(1-7) , but late recovery is enhanced compared to filgrastim.

-AII(1-7) improves the recovery towards baseline in all formed blood elements and minimizes pre-cycle progressive pancytopenia associated with myelosuppressive chemotherapy.

It is to be understood that the invention is not to be limited to the exact details of operation, or to the exact compounds, compositions, methods, procedures or embodiments shown and described, as obvious modifications and equivalents will be apparent to one skilled in the art, and the invention is therefore to be limited only by the full scope of the appended claims.

We claim:

1. An improved method for chemotherapy in a human patient, wherein the improvement comprises administering to the human chemotherapy patient an amount effective for treating or preventing chemotherapy side effects of at least one active agent comprising a sequence of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



wherein  $R^1$  is selected from the group consisting of H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^2$  is selected from the group consisting of Arg, Lys, Ala, Orn, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, Lys, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ser, Ala, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is selected from the group consisting of His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is selected from the group consisting of Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr,

excluding sequences including  $R^4$  as a terminal Tyr group;

and wherein the active agent is not SEQ ID NO:1.

2. The method of claim 1 wherein the active agent comprises a sequence of at least four contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

DocId: 267260

3. The method of claim 1 wherein the active agent comprises a sequence of at least five contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

4. The method of claim 1 wherein the active agent comprises a sequence of at least six contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

5. The method of claim 1 wherein the active agent comprises a sequence of at least seven contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

6. The method of claim 1 wherein the active agent consists essentially of a sequence of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

7. The method of claim 1 wherein the active agent consists essentially of a sequence of at least four contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

8. The method of claim 1 wherein the active agent consists essentially of a sequence of at least five contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

9. The method of claim 1 wherein the active agent consists essentially of a sequence of at least six contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

10. The method of claim 1 wherein the active agent consists essentially of a sequence of at least seven contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I.

11. The method of claim 1 wherein the active agent comprises a sequence selected from the group consisting of angiotensinogen, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO: 34; SEQ ID NO:35, SEQ

ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, and SEQ ID NO:42.

12. The method of claim 1 wherein the active agent comprises the amino acid sequence of SEQ ID NO:4.

5 13. The method of claim 1 wherein the active agent consists essentially of a sequence selected from the group consisting of angiotensinogen, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, 10 SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO: 34; SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, and SEQ ID NO:42.

14. The method of claim 1 wherein the active agent consists essentially of the amino acid 15 sequence of SEQ ID NO:4.

15. The method of claim 1 wherein the active agent comprises a sequence of the following general formula II:

Asp-Arg-R1-R2-Ile-His-Pro-R3, wherein

R1 is selected from the group consisting of Val, Pro, Lys, Norleu, and Leu;

20 R2 is selected from the group consisting of Ala, Tyr, and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe or is absent.

16. The method of claim 1 wherein the active agent consists essentially of a sequence of the following general formula II:

Asp-Arg-R1-R2-Ile-His-Pro-R3, wherein

R1 is selected from the group consisting of Val, Pro, Lys, Norleu, and Leu;

R2 is selected from the group consisting of Ala, Tyr, and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe or is absent.

17. The method of claim 15 wherein the active agent comprises a sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, and SEQ ID NO:41.

18. The method of claim 16 wherein the active agent consists essentially of a sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, and SEQ ID NO:41.

19. The method of claim 1 wherein the side effect is selected from the group consisting of hematopoietic toxicity, decreased mobilization of hematopoietic progenitor cells from bone marrow into the peripheral blood; anemia, myelosuppression, pancytopenia, thrombocytopenia, neutropenia, lymphopenia, leukopenia, stomatitis, alopecia, headache, and muscle pain.

20. The method of claim 1 wherein the active agent is administered at a dosage of between about 2.5 µg/kg/day and about 100 µg/kg/day.

21. The method of claim 1 wherein the active agent is administered at a dosage of between about 10 µg/kg/day and about 75 µg/kg/day.

22. The method of claim 1 wherein the active agent is administered parenterally.

23. The method of claim 22 wherein the active agent is administered subcutaneously or intravenously.

24. The method of claim 23 wherein the active agent is self-administered.

25. The method of claim 24 wherein the active agent is administered into the abdomen or thigh.

26. The method of claim 1 wherein administration of the active agent is initiated either at the time chemotherapy is initiated, or subsequently to initiation of chemotherapy.

27. The method of claim 1 wherein the active agent is administered once per day.

28. A pharmaceutical composition comprising

a) an amount of the active agent of claim 1 sufficient to provide a dosage to a patient of between about 2.5 µg/kg/day and about 100 µg/kg/day; and

b) a pharmaceutically acceptable carrier.

29. The pharmaceutical composition of claim 28 wherein the active agent is selected from the group consisting of SEQ ID NO:4 and SEQ ID NO:41

30. The pharmaceutical composition of claim 28 further comprising an amount effective of a cytokine for increasing hematopoietic cell production.

31. The pharmaceutical composition of claim 30 wherein the cytokine is selected from the group consisting of granulocyte colony stimulating factor, granulocyte-macrophage-CSF, epidermal growth factor, interleukin 11, thrombopoietin, megakaryocyte development and growth factor, pixykin, stem cell factor, FLT-ligand, and interleukins 1, 3, 6, and 7.

32. The pharmaceutical composition of claim 31 wherein the cytokine is granulocyte colony stimulating factor.

33. An article of manufacture, comprising the pharmaceutical composition of claim 28 loaded in a drug delivery device.

34. The article of manufacture of claim 33 wherein the delivery device is a syringe.



## Abstract of the Disclosure

The present invention provides improved methods, kits, and pharmaceutical compositions for increasing hematopoietic cell survival and/or reducing or preventing the side effects of chemotherapy, and mobilizing hematopoietic progenitor cells from bone marrow into peripheral  
5 blood following chemotherapy, comprising administering an effective amount of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists to a patient in need of chemotherapy.

002277-267E2260

# Effect of Angiotensin II on White Blood Cell Number





Fig 1

Blood, day 7.

000000 26 FEB 2000

# Effect of Angiotensin II on White Blood Cell Number



Treatment Group

Spleen, day 7.

0002277-267E2650

# Effect of Angiotensin II on White Blood Cell Number





Fig 3

Thymus, day 7



BM, day 7.

# Effect of Angiotensin II on CFU-GM (Day 7)




Fig. 5

Blood, day 7.

002277-267E2450

# Effect of Angiotensin II on CFU-GM (Day 7)





Fig 6

Spleen, day 7.

# Effect of Angiotensin II on CFU-GM (Day 7)



BM, day 7.



# Effect of Angiotensin II on CFU-GM




Fig 8

Blood, day 7.

# Effect of Angiotensin II on White Blood Cell Number





Fig 9


Spleen, day 14

# Effect of Angiotensin II on White Blood Cell Number




Thymus, day 14

# Effect of Angiotensin II on White Blood Cell Number




BM, day 14

# Effect of Angiotensin II on CFU-GM (Day 7)



Spleen, day 14

# Effect of Angiotensin II on CFU-GM (Day 7)



Blood, day 14

# Effect of Angiotensin II on CFU-GM (Day 7)




Fig 14

Blood, day 14

# Effect of Angiotensin II on CFU-GM (Day 7)





Fig 15

Spleen, day 14



0022277 26722250

# Effect of Angiotensin II on CFU-GM (Day 7)



# Effect of Angiotensin II on CFU-GM (Day 7)




Fig. 17

Femur, day 14

# Effect of Angiotensin II on CFU-GM




Fig 18

BM, day 7.

# Effect of Angiotensin II on CFU-GM





Fig 19

Spleen, day 7.

# Effect of Angiotensin II on White Blood Cell Number



Blood, day 14.

# Effect of AI on WBC after Chemotherapy





Fig 71

002277-2672260

# Effect of All(1-7) on WBC after Chemotherapy



# Effect of 1GD on WBC after Chemotherapy





Fig 23




004217 46FE2260

# Effect of 2GD on WBC after Chemotherapy



# Effect of 5GD on WBC after Chemotherapy



# Effect of 9GD on WBC after Chemotherapy





Fig 2b


# Effect of Angiotensin Peptides on GM-CFU



Chemotherapy, Bone Marrow, 10 ug, Day 10

Fig 27


# Effect of Angiotensin Peptides on GM-CFU



Chemotherapy, Bone Marrow, 100 ug, Day 10

Fl<sub>15</sub><sup>28</sup>

# Effect of Angiotensin Peptides on GM-CFU



Chemotherapy, Blood, 10 ug

# Effect of Angiotensin Peptides on GM-CFU




Fig 30  
30

Chemotherapy, Blood, 100 ug

Figure 31A. Effect of A(1-7) on Hematopoietic Recovery after Cytosin Chemotherapy




Figure 31B. Effect of A(1-7) on Hematopoietic Recovery after Cytosin Chemotherapy






Figure 3A. Effect of A(1-7) on Hematopoietic Recovery after Cytosin Chemotherapy




Figure 3B. Effect of A(1-7) on Hematopoietic Recovery after Cytosin Chemotherapy




Figure 33 Effect of Angiotensin Fragments on WBC




Figure 34 Effect of Angiotensin Fragments on WBC




Figure 35 Effect of Angiotensin Fragments on WBC




Figure 36 Effect of Angiotensin Fragments on Platelets




Figure 37 Effect of Angiotensin Fragments on Platelets




Figure 38 Effect of Angiotensin Fragments on Platelets




Figure 39 Effect of Angiotensin Fragments on Hemoglobin




Figure 40 Effect of Angiotensin Fragments on Hemoglobin




Figure 4 | Effect of Angiotensin Fragments on Hemoglobin




Figure 1 consists of 12 line graphs, labeled (a) through (l), showing the evolution of the number of species in different taxonomic groups over time. The groups are: (a) All species, (b) Invertebrates, (c) Vertebrates, (d) Fish, (e) Amphibians, (f) Reptiles, (g) Birds, (h) Mammals, (i) Insects, (j) Arachnids, (k) Molluscs, and (l) Plants. Each graph plots the number of species (Y-axis) against time in millions of years (X-axis). The data points are connected by lines, showing the growth of species over time. The Y-axis scale varies for each group, and the X-axis scale also varies. The graphs show that the number of species in all groups increases over time, with a significant increase in the number of species in the vertebrate groups (d, e, f, g, h) starting around 200 million years ago.

<210> 4

<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (1-7)

<400> 4  
Asp Arg Val Tyr Ile His Pro  
1 5

<210> 5  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (2-7)

<400> 5  
Arg Val Tyr Ile His Pro  
1 5

<210> 6  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (3-7)

<400> 6  
Val Tyr Ile His Pro  
1 5

<210> 7  
<211> 4  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (5-8)

<400> 7  
Ile His Pro Phe  
1

<210> 8  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>



<223> Description of Artificial Sequence: AII (1-6)

<400> 8

Asp Arg Val Tyr Ile His  
1 5

<210> 9

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (1-5)

<400> 9

Asp Arg Val Tyr Ile  
1 5

<210> 10

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (1-4)

<400> 10

Asp Arg Val Tyr  
1

<210> 11

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (1-3)

<400> 11

Asp Arg Val  
1

<210> 12

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue

<220>

<221> MOD\_RES

<222> (2)

<223> Nle

<400> 12

Arg Xaa Tyr Ile His Pro Phe

1

5

<210> 13

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue

<220>

<221> MOD\_RES

<222> (4)

<223> Nle

<400> 13

Arg Val Tyr Xaa His Pro Phe

1

5

<210> 14

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (6-8)

<400> 14

His Pro Phe

1

<210> 15

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (4-8)

<400> 15

Tyr Ile His Pro Phe

1

5

<210> 16

<211> 7

<212> PRT

<213> Artificial Sequence

<220>



<220>  
<223> Description of Artificial Sequence: AII analogue 1

<400> 19  
Asp Arg Val Tyr Val His Pro Phe  
1 5

<210> 20  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 2

<400> 20  
Asn Arg Val Tyr Val His Pro Phe  
1 5

<210> 21  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 3

<400> 21  
Ala Pro Gly Asp Arg Ile Tyr Val His Pro Phe  
1 5 10

<210> 22  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 4

<400> 22  
Glu Arg Val Tyr Ile His Pro Phe  
1 5

<210> 23  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 5

<400> 23  
Asp Lys Val Tyr Ile His Pro Phe

004247-26250

1

5

<210> 24

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 6

<400> 24

Asp Arg Ala Tyr Ile His Pro Phe

1

5

<210> 25

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 7

<400> 25

Asp Arg Val Thr Ile His Pro Phe

1

5

<210> 26

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 8

<400> 26

Asp Arg Val Tyr Leu His Pro Phe

1

5

<210> 27

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 9

<400> 27

Asp Arg Val Tyr Ile Arg Pro Phe

1

5

<210> 28

<211> 8

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

```
<400> 28
Asp Arg Val Tyr Ile His Ala Phe
  1             5
```

<220>  
<223> Description of Artificial Sequence:All analogue 11

```
<400> 29
Asp Arg Val Tyr Ile His Pro Tyr
  1             5
```

```
<210> 30
<211> 8
<212> PRT
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence: AII analogue 12

```
<400> 30
Pro Arg Val Tyr Ile His Pro Phe
      1             5
```

```
<210> 31
<211> 8
<212> PRT
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence: AII analogue 13

```
<400> 31
Asp Arg Pro Tyr Ile His Pro Phe
  1                               5
```

```
<210> 32
<211> 8
<212> PRT
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence: AII analogue 14



[illegible]

```
<210> 39
<211> 7
<212> PRT
```



Figure 1 consists of a 3x3 grid of bar charts. Each chart has a vertical axis labeled 'Percentage' ranging from 0 to 100. The horizontal axis for each chart lists nine categories: 1. No answer, 2. No answer, 3. No answer, 4. No answer, 5. No answer, 6. No answer, 7. No answer, 8. No answer, 9. No answer. The bars are colored in shades of gray. The charts show varying percentages for each category across the three groups: 'All respondents', 'Non-respondents', and 'Respondents'.

<223> Description of Artificial Sequence:2GD:  
Pro3-AII(1-7)

Asp Arg Pro Tyr Ile His Pro  
1 5

<213> Artificial Sequence

<223> Description of Artificial Sequence:5GD:  
Lys3-AII(1-7)

Asp Arg Lys Tyr Ile His Pro  
1 5

<213> Artificial Sequence

<223> Nle

<223> Description of Artificial Sequence:9GD:  
noreu-AII(1-7)

Asp Arg Xaa Tyr Ile His Pro  
1 5